


<input checked="" type="checkbox"/> APPROVED FOR DISTRIBUTION
 (APPROVING OFFICIAL)
12/15/83 (DATE)

Reproduction of Weakfish, Cynoscion regalis, in the New York Bight  
and Evidence for Geographically Specific  
Life History Characteristics

by

Gary R. Shepherd  
Rutgers University  
New Brunswick, New Jersey 08903<sup>1</sup>

and

Churchill B. Grimes  
Rutgers University  
New Brunswick, New Jersey 08903

<sup>1</sup>Present address:  
National Marine Fisheries Service  
Northeast Fisheries Center  
Woods Hole Laboratory  
Woods Hole, Massachusetts 02543

## ABSTRACT

Reproduction characteristics for weakfish, Cynoscion regalis, in the New York Bight were examined. Spawning in 1980-81 occurred from May to early July with spawning time dependent on parental size. Maturity for both sexes occurred by age one but at a greater size in females. Annual fecundity estimates were compared to literature values for North Carolina weakfish and were found to be considerably lower at size, yet cumulative fecundities were nearly equivalent. The latitudinal variations in fecundity may be a behaviorally and environmentally induced phenomena, and influence the long term population stability of weakfish.

## INTRODUCTION

Weakfish, Cynoscion regalis, are a member of the family Sciaenidae and are a common inshore species occurring between Cape Cod, Massachusetts, and southern Florida. The species undergoes a spring migration from offshore waters of Virginia and the Carolinas to appropriate estuarine spawning areas, then a return migration in late fall to overwintering grounds (Nesbit 1954). The center of greatest abundance occurs within the Middle Atlantic Bight in quantities sufficient to support a recreational and commercial fishery. In 1979 commercial fishermen landed 13,000 metric tons (mt) of weakfish and nearly 5,000 mt were caught by recreational anglers (Wilk 1981). Abundance has, however, undergone some dramatic fluctuations over the last several decades. Commercial landings averaged 8,800 mt from 1940-49, then dropped to 2,915 mt by 1950 and remained at these low levels until the mid-1970's (Wilk 1981). The exact cause of these variations remains a mystery, although speculations include over-fishing, DDT-induced mortality, and environmentally induced recruitment failure (Joseph 1972, Massman 1963, Merriner 1976). To adequately assess the mechanisms controlling recruitment success or failure, we must first have a thorough understanding of the reproductive biology of weakfish.

Merriner (1976) has examined reproduction of weakfish in North Carolina, and Daiber (1957) mentioned spawning behavior of weakfish in Delaware Bay, but the reproductive biology of weakfish in their northern range has never been fully investigated. Furthermore, there is reason to believe that reproduction may vary throughout the geographic range. Leggett and Carscadden (1978) have shown latitudinal variations in reproduction and growth of American shad (Alosa sapidissima) and White and Chittenden (1977) have likewise shown

geographic differences in another sciaenid, the Atlantic croaker (Micropogonias undulatus). Growth differences have already been established for weakfish between the New York Bight and North Carolina (Shepherd and Grimes<sup>2</sup>, Perlmutter, Miller and Poole 1956), so there is reason to suspect possible reproductive differences. The purpose of the study was to investigate weakfish reproduction in the Middle Atlantic region, determine if any geographic variations exist and consider possible reasons for geographically specific characteristics.

## METHODS AND MATERIALS

### Sample Collection

Samples (n=1208) were collected during the National Marine Fisheries Service groundfish survey from 1980-83 at stratified random stations north of Chesapeake Bay (Fig. 1). Fish were collected with a #41 Yankee trawl in spring and a #30 Yankee trawl in summer and fall at depths between 5 and 200 m (Grosslein 1969). NMFS samples were supplemented by 461 fish collected between May 1980 and June 1981 from commercial pound nets in Gardiners Bay, New York, (n=61) and Sandy Hook Bay, New Jersey, (n=115) and from pair and otter trawl fisheries in Delaware Bay (n=285).

At each NMFS station, weakfish catches were stratified into 10 cm size intervals and 10-15 fish sampled per interval. Total length (TL) to the nearest cm, sex, and maturity stage were recorded. Weakfish from Gardiners Bay were randomly sampled, TL measured to the nearest mm, sex, and maturity stage recorded, and gonads removed and weighed to the nearest 0.1 g. Sandy Hook and Delaware Bay samples were collected by random selection of 50 lb (22.5 kg) boxes in each market category available from the catch and TL to the

nearest mm, and weight (whole and gutted) to the nearest g recorded. Gonads and livers were also removed and weighed to the nearest 0.1 g. Gonads were preserved in modified Gilson's solution (Bagenal and Braum 1971) for several weeks, then removed, washed with distilled water and stored in 95% isopropyl alcohol.

#### Maturity and Fecundity Methods

The maturity stage of each sample was categorized as immature, developing, ripe (spawning), spent, or recovering as modified from Nikolsky (1963). The maturity stages were further sub-divided as mature or immature for calculation of length at 50% maturity using probit analysis (Finney 1971).

Seasonality of reproduction was determined from changes in the gonad condition. A gonosomatic index (GSI) was calculated to show changes in gonad weight relative to somatic weight (gutted body weight). The index was computed as:

$$\text{GSI} = (\text{gonad weight}/\text{somatic weight}) \times 100$$

To examine physiological changes associated with spawning, liver condition was assessed using a hepatosomatic index (HSI) computed by substituting liver weight for gonad weight in the above relationship (Hut-Han 1978). The differences in monthly mean hepatosomatic indices between sexes were analyzed statistically using a Wilcoxon test (Patzner and Adam 1981).

The number of spawnings in the season were investigated by analyzing the seasonal frequency distribution of oocyte diameters (Hickling and Rutenberg 1936). In each of 15 samples collected between 5 May and 22 July 1980-81 in Delaware and Gardiners Bays, three subsamples were taken per ovary and approximately 500 oocyte diameters were randomly measured using an ocular micrometer.

Fecundity estimates were determined from 28 fish macroscopically classified as developing, which were captured during May 1981 in Delaware and Gardiners Bays. An oocyte diameter of 0.20 mm was determined from diameter frequency distributions and the degree of yolk accumulation as the size between oogonia and developing ova, and was used as the lower size limit of ova in the fecundity estimates. Each sample was diluted with distilled water, stirred, then several aliquots removed from the solution to provide a density of approximately 10,000 ova in a 6 cm x 6 cm gridded petri dish. About 800-1000 ova were counted from six randomly selected squares, then adjusted for a total sub-sample count. Two sub-samples were counted per sample and three if the ova sample was from a large fish (>60 cm). The sample and subsample were oven dried at 40°C for a minimum of 24 hours then weighed to the nearest 0.001 g on a Mettler balance. Total fecundity was calculated as:

$$\text{Total fecundity} = [\text{number in subsample} \times (\text{sample wt} / \text{subsample wt})] + \# \text{ in subsample}$$

Predictive equations of fecundity from length and weight were fit to a geometric mean (GM) functional regression (Ricker 1975) following log-log transformation.

## RESULTS

### Seasonality

The changes in maturity stages during the year indicate spawning takes place from May to mid-July for weakfish in Delaware Bay and north to Long Island (Fig. 2). In May, all 122 mature fish examined of both sexes were in the developing or ripe stage of gonad development. The 50 fish that were inspected in June included 100% of the males and 70% of the females in a developing or ripe stage. Several of the ovaries examined in a June sample

from Gardiners Bay had the appearance of being partially spent. The ovary was flacid, slightly hemorrhaged and the lumen filled with fluid, but a few transparent ova were still visible. Spawning weakfish in Delaware Bay were captured as late as 12 July, when 11% of the females and 76% of the males were classified as ripe. In the same month, 84% of the females were in spent condition. By August, all of the fish examined at all locations were in post-spawning condition. Weakfish from Sandy Hook Bay were not available until June and all the samples examined from this area were in post-spawning condition, either spent or recovering.

This seasonality of spawning at each of the sample locations was suggested by seasonal changes in gonasomatic indices (GSI). Mean GSI values (% gonad weight relative to somatic weight)  $\pm$  95% CI of all females rose from  $5.75 \pm 1.29$  in May to a peak value of  $6.04 \pm 1.43$  in June, then declined to  $1.76 \pm 0.40$  by July (Fig. 3). Male gonosomatic indices were at a yearly high of  $4.88 \pm 1.45$  in May but only declined to  $2.51 \pm 0.85$  by July. The mean GSI values for females were consistently higher in all months except July. Gonad size reached the lowest levels for both sexes in August with mean GSI values of  $0.71 \pm 0.07$  and  $0.19 \pm 0.02$  for females and males, respectively. The GSI values remained low until the last samples were collected in November.

Specific spawning time, as reflected in gonasomatic indices, was dependent on the size of the individual fish (Fig. 4). Mean index values and lengths  $\pm$  95% CI for developing fish in Delaware and Gardiners Bays during May was  $2.48 \pm 0.67$  at a mean length of  $551 \text{ mm} \pm 47.0$  for females and  $1.13 \pm 0.85$  at a length of  $391 \text{ mm} \pm 98.9$  for males. Spawning fish captured during May had indices of  $8.65 \pm 1.64$  at mean length of  $702 \text{ mm} \pm 25.2$  for females and  $6.49 \pm 1.58$  at mean length of  $619 \text{ mm} \pm 33.1$  for males. In June, spawning fish had mean GSI values of  $6.72 \pm 1.83$  with mean length of  $544 \text{ mm} \pm 95.6$  for females

and  $4.54 \pm 1.04$  with mean length of  $570 \text{ mm} \pm 56.3$  for males. These index variations confirmed observations of earlier spawning by large fish.

Seasonal variation in mean hepatosomatic indices (HSI) of Delaware and Sandy Hook Bay samples indicates that changes in liver weight occurred at the time of spawning and prior to fall migration (Fig. 5). Mean HSI values  $\pm$  95% CI for females decreased from a high in May of  $2.80 \pm 0.35$  to a low in September of  $0.87 \pm 0.30$ . The index values for males followed the same pattern declining from  $1.89 \pm 0.27$  in May to  $0.96 \pm 0.18$  in September, but the maximum values occurred in November with a mean HSI of  $2.57 \pm 0.29$ . The indices were tested for differences between sexes and the differences were found to be significant in May and June ( $P < 0.001$ ) with the values for females being greater (Table 1). No significant differences were found in a comparison of the July through November samples.

Frequency distributions of oocyte diameters from 15 fish, ranging from 55 to 81 cm and collected between 6 May and 22 July show one seasonally progressing mode of developing ova (Fig. 6). The position of the mode varied according to the development stage of the individual ovary. A sample from 5 May had a mode between 0.02 and 0.45 mm and was skewed toward a prominent peak at 0.12 mm. In the 13 May sample, a second mode appeared around 0.35 mm and this mode increased to a size of 0.63 mm in the 20 June sample. The 22 July sample contained only oogonia. Maximum ova diameter observed was 0.95 mm from a sample on 26 May. That ovum was filled with fluid between the yolk and chorion, indicative of an ova immediately prior to release (Bagenal 1967). This corresponds to the size of weakfish eggs, 0.870 to 0.975 mm, which have been identified in the water column (Harmic 1958).

### Maturity, Sex Ratio, and Fecundity

Length  $\pm$  95% CI at which 50% of the total sampled population reached maturity was similar for both sexes. Females attained 50% maturity at 25.6 cm  $\pm$  1.2 cm while males were slightly lower at 25.1 cm  $\pm$  1.1 cm (Table 2). The corresponding age at maturity for both sexes was one year. The smallest mature male and female was 20 cm. The maximum size of immature weakfish was 40 cm for females and 33 cm for males.

The overall population sex ratio approached equality (Table 3). The sex ratio of the population, divided into 5 cm length intervals, was 48 to 52 females to males and was not significantly different from 50:50 as determined from a chi-square analysis ( $\chi^2 = 1.81$ ,  $\chi^2_{0.05} = 3.84$ ,  $n = 1669$ ). Sex ratio at size data did reveal significant differences from 50:50 for several length intervals. At 40 cm, the ratio was 52:48 female to male but increased to 60.5% ( $\chi^2 = 5.25$ ,  $\chi^2_{0.05} = 3.84$ ,  $n = 119$ ) and 63.8% ( $\chi^2 = 7.19$ ,  $\chi^2_{0.01} = 6.64$ ,  $n = 94$ ) males at 45 and 55 cm, respectively. The sex ratio of combined length intervals above 55 cm had a ratio of 58.5:41.5 which was significantly dominated by females ( $\chi^2 = 7.45$ ,  $\chi^2_{0.01} = 6.64$ ,  $n = 260$ ).

Regression models indicated that length and weight were equally predictive of fecundity. The relationships between total length or gutted weight, and fecundity were best described by a power curve. The GM functional regression (Ricker 1975) of the  $\log_e$  transformed data  $\pm$  standard error of the regression were:

$$\ln \text{ fecundity} = -16.322 + 4.659 \ln \text{ total length (mm)} \pm 0.368$$

$$r^2 = 0.835 \quad n = 28$$

Fig. 7

$$\ln \text{ fecundity} = 1.975 + 1.542 \ln \text{ gutted weight (g)} \pm 0.364$$

$$r^2 = 0.839 \quad n = 28$$

Fecundities were also estimated for six weakfish collected in June from Gardiners Bay which appeared to be spent, although they had ova remaining in the gonad. These ova were 0.55 to 0.65 mm which were similar in size to those present in a ripe ovary. All six samples were significantly different ( $P < 0.001$ ) than expected values based on a Student's t-test (Table 4). These June samples were 60-75% lower than the predicted fecundities at length.

#### DISCUSSION

Weakfish spawning in northern estuaries is a seasonal event which occurs following the spring inshore migration (Hildebrand and Schroeder 1928, Welsh and Breder 1923). Our study found the spawning period for weakfish in Delaware Bay and Gardiners Bay, as determined from maturity stages, to be from May to mid-July. Ichthyoplankton surveys have found weakfish eggs and larvae present in New York Bight from May to July (Colton et al. 1979). The spawning period was further substantiated by changes in gonosomatic indices (GSI). The mean GSI values reached a maximum in May for males and in June for females, then both declined to minimum values by August. The gonad weights were maintained at this low level until the fish disappeared from the coast in late November. The spawning period of weakfish in North Carolina is somewhat longer, extending from March to September with the peak period from April to June (Merriner 1976). The duration of the ripe maturity stage in Delaware and Gardiners Bays were greater for males, with ripe males captured as late as 22 July. The protracted spawning season of males was also evident in the GSI values. July was the only month in which male indices were greater than those of females, indicating that the gonad weight per unit body weight was larger

in males. In other months, female GSI values were as much as 2.8% greater than males.

Sex differences were also evident in physiological changes associated with spawning. HSI values were near maximum in May, during spawning season, and as expected higher in females. Development of ovaries demands higher energetic costs than testicular development (Nikolsky 1963), therefore the necessary energy reserves in the liver would be proportionately larger in females (Timashova 1982). After spawning the sexual dimorphism disappears and HSI values for both sexes increase similarly prior to the fall migration.

The peak spawning period in the New York Bight estuaries varied with different size weakfish. Generally, among migratory fishes, the larger individuals will return to an estuary prior to their smaller counterparts (Briggs 1955, Nikolsky 1963). Weakfish exhibit a similar behavior, as the largest individuals or "tiderunners" enter the bays in early May and spawn by mid-May whereas the smaller weakfish, arrive later and reach peak spawning during June. The GSI values for fish greater than 60 cm generally decline from May to June, while smaller fish increase to maximum values in June. This differential spawning based on parental size results in two spawning peaks and the subsequent appearance of bimodal length frequency data for juvenile weakfish (Shepherd and Grimes<sup>3</sup>, Daiber 1954, Thomas 1971).

#### Sex Ratio, Maturity, and Fecundity

The overall sex ratio of the population is not different from 1:1, but one sex was dominant at certain length intervals. We believe the deviations from a 1:1 ratio at various lengths was attributable to differential growth between sexes (Wenner 1972). Female growth begins to exceed male growth at

approximately 45-55 cm (Shepherd and Grimes<sup>4</sup>), at which point the sex ratio becomes dominated by males. Females grew beyond that size interval faster and occupied the majority of the 60-85 cm length intervals. Although males have growth potential equal to females, the numbers attaining maximum size were greater for females.

In northern waters, the size at which weakfish attained 50% population maturity was similar for both sexes. Females matured by 25.6 cm and age 1 while males attained maturity at 25.1 cm, also at age 1. Apparently, greater differences exist between northern weakfish (Delaware Bay and north) and southern weakfish (North Carolina). Although ages were similar, southern females spawned at 23 cm and males by 18 cm (Merriner 1973).

Estimates of fecundity for New York Bight weakfish differ from estimates for southern weakfish. Weakfish in North Carolina did not reach sizes much beyond 45 cm but had fecundities, relative to length, several times higher than northern weakfish (Merriner 1976) (Table 5). For example, at 50 cm TL female weakfish from New York Bight produced 306,159 ova while the fecundity of southern fish of the same size was 2,051,080 ova. In spite of these large differences, lifetime fecundity would be approximately equal. Southern weakfish can potentially reproduce until age 5 and produce approximately 9,913,080 ova (based on the equation  $\text{fecundity} = 0.152 \text{ TL}^{2.6418}$  from Merriner 1976), whereas northern weakfish reproducing for 10 years have nearly equivalent total ova production of 10,008,167.

Batch spawning, involving two distinct groups of ova, was found for weakfish in North Carolina (Merriner 1976). In samples examined in 1980-81 from Delaware and Gardiners Bays, multiple spawns were not evident. The ova diameter frequencies of developing and ripe ovaries contained two modes, one

consisting of reserve oocytes and another of developing ova. The developing ova had a wide size range ( $\approx 0.3$  mm) and may have been released during consecutive spawning events, but did not constitute separate batches within an ovary. Furthermore, all ova produced annually by weakfish may not be released during spawning. A study of Delaware Bay weakfish in 1954 proposed batch spawning, but spent ovaries were not examined to determine if all ova were released (Daiber 1954). Ovaries classified as spent which we examined still contained 25-40% of the ova expected for a fish of that size. These results suggest fertility may be 60-75% of estimated potential fecundity. Foucher and Beamish (1980) reported similar conclusions from studies of Pacific hake (Merluccius productus). We did not examine enough spent weakfish ovaries to determine if this phenomenon was consistent from year to year.

The variable reproductive and age and growth characteristics for weakfish in different geographic areas suggest specific physiological responses to different environmental conditions. North Carolina weakfish had smaller sizes at maturity, smaller length at corresponding age after age 1, reduced longevity and maximum size, and higher relative fecundity than New York Bight weakfish. However, the lifetime reproductive potential was nearly equal for both groups. These life history characteristics for weakfish are similar to clinal variations between Labrador and Florida described for American shad (Alosa sapidissima) (Leggett and Carscadden 1978), differences in reproductive characteristics between Atlantic herring (Clupea harengus) in the Norway and the Baltic Sea (Schopka 1971), and north to south variations in North American populations of Pacific herring (Paulson and Smith 1977).

The different reproductive strategies in weakfish may result from varying environmental demands. When weakfish spawning occurs in northern estuaries

water temperatures are unpredictable and subject to sudden drops which are potentially lethal to eggs and larvae (Harmic 1958). Table 6 shows minimum estuarine temperatures during the spawning season in northern waters may drop below the temperature limits of 12-16°C necessary for successful hatching (Harmic 1958). The higher probability of pre-reproductive mortality of progeny in northern estuaries results in a "bet-hedging" strategy in which fewer eggs are produced each year, but the possible number of annual spawnings are increased (Stearns 1976, Giesel 1976), thus maximizing potential contributions to the gene pool throughout a fish's lifespan. In contrast, southern weakfish spawn in a more predictable estuarine environment (Table 4) and consequently there is less chance of environmentally induced pre-reproductive mortality. However, southern fish are faced with greater post-reproductive mortality (longevity observed by Merriner (1973) was 5 years). Greatest reproductive success in this situation requires maximizing annual gamete production in the few years possible. In addition, weakfish migrating to northern estuaries (Nesbit 1954) may utilize energy reserves otherwise available for gonad growth, whereas southern fish having less distance to travel may reallocate energy towards reproduction.

Consequences of the area specific reproductive characteristics may be a reduced population stability for weakfish in the northern end of the range. Apparently, northern fish have a strategy to cope with potentially higher egg and larval mortality by spreading reproduction over more years and reducing annual fecundity, i.e., a "bet-hedging" strategy (Stearns 1976). Therefore, to fulfill their reproductive potential they must avoid premature adult mortality. If adult mortality (natural and fishing) in weakfish becomes excessive, the larger, most fecund individuals will be lost or reduced, thus shifting the burden of spawning to the smaller, less fecund fish. When the

value of  $b$  in the fecundity equation  $F = aTL^b$  is greater than 3, as in New York Bight weakfish, then truncation of the size/age structure in a spawning population will also result in a reduction of population fecundity (Hempel 1979). Concurrent high adult mortality and high pre-reproductive mortality could contribute to a decline in population abundance. The large fluctuations which have occurred in weakfish populations over the last several decades (Wilk 1979) may be due in part to these circumstances (i.e., high adult and pre-reproductive mortality). Although the correlations between population fecundities and recruitment is not usually strong for marine fishes (Cushing 1977), a decrease in population fecundity may eventually reduce any buffer that weakfish have of withstanding natural fluctuations in egg, larval, and juvenile survival. Therefore, if management practices are to effectively regulate the weakfish resources, geographic variations in reproductive potential should be considered.

## ACKNOWLEDGMENTS

The authors would like to thank Drs. Ken Able and J. Richard Trout for their helpful guidance. Mr. Dan Dzenkowski and crew of the Dan Dee, Greenport, L.I., were more than generous in allowing us to sample their catches and personnel of NMFS, Woods Hole, MA, especially Dr. Ambrose Jearld, kindly provided samples from the groundfish surveys. We would also like to thank Steve Turner, Chuck Idelberger and Susan Toogood for their assistance and Hannah Goodale for her assistance in preparing the manuscript. Funding for this project provided by NMFS (contract #NA-79-FAC-00041) and the New Jersey Agricultural Experiment Station (NJAES publication #D-12409-14-83).

## LITERATURE CITED

- Bagenal, T.B. 1967. A short review of fish fecundity. In: S.D. Gerking (ed.), The Biological Basis of Freshwater Fish Production. John-Wiley & Sons, N.Y. 496 p.
- Bagenal, T.B. and E. Braum. 1971. Eggs and early life history. In: Ricker, W.E. (ed.), Methods for Assessment of Fish Production in Fresh Waters. Blackwell Sci. Publ., Oxford, Eng., 365 p.
- Briggs, J.C. 1955. Behavior pattern in migratory fishes. Science 122 (3162): 240.
- Colton, J.B., Jr., A.W. Smith, W. Kendall, Jr., P.L. Berrien, and M.P. Fahay. 1979. Principal spawning areas and times of marine fishes, Cape Sable to Cape Hatteras. Fish. Bull. 76(4): 911-915.
- Cushing, D.H. 1974. The natural regulation of fish productions. In: Harden-Jones, F.R. (ed.), Sea Fisheries Research, John-Wiley and Sons, London, England, 510 p.
- Daiber, F.C. 1954. Fisheries research program. Mar. Lab. Dep. Biol. Sci., Univ. Del. Biennial Rep. 1958 and 1954. Publ. 2:50-64.
- Daiber, F.C. 1957. Sea trout research. Estuarine Bull. 2(5): 1-6.
- Finney, D.J. 1971. Probit Analysis. Cambridge Univ. Press, Cambridge. 333 p.
- Foucher, R.P. and R.J. Beamish. 1980. Production of nonviable oocytes by Pacific hake (Merluccius productus). Can. J. Fish. Aquat. Sci. 37: 41-48.
- Giesel, J.T. 1976. Reproductive strategies as adaptations to life in temporarily heterogeneous environments. Ann. Rev. Ecol. Syst. 7: 57-70.

- Grosslein, M.D. 1969. Groundfish survey programs of BCF Woods Hole. Commer. Fish. Rev. 31(8-9): 22-35.
- Harmic, J.L. 1958. Some aspects of the development and ecology of the pelagic phase of the grey squeteague Cynoscion regalis (Block and Schneider), in the Delaware estuary. Ph.D. Thesis, Univ. Delaware, Newark, Del. 167 p.
- Hempel, G. 1979. Early Life History of Marine Fish: The Egg Stage. Univ. Washington Press. 70 pp.
- Hickling, C.F. and E. Rutenberg. 1936. The ovary as an indicator of the spawning period in fishes. J. Mar. Biol. Assoc. U.K. 21: 311-316.
- Hildebrand, S.F. and W.C. Schroeder. 1928. Fishes of Chesapeake Bay. U.S. Bureau of Fisheries 43(1): 1-366.
- Hut-Han, M. 1978. The reproductive biology of the dab, Limanda limanda (L.) in the North Sea: gonosomatic index, hepatosomatic index and condition factor. J. Fish. Biol. 13: 369-378.
- Joseph, E.B. 1972. The status of the sciaenid stocks of the Middle Atlantic coast. Chesapeake Sci. 13(2): 87-100.
- Leggett, W.C. and J.E. Carscadden. 1978. Latitudinal variation in reproductive characteristics of American shad (Alosa sapidissima); evidence for population specific life history strategies in fish. J. Fish. Res. Bd. Can. 35: 1469-1478.
- Massman, W.H. 1963. Age and size composition of weakfish, Cynoscion regalis, from pound nets in Chesapeake Bay, Virginia 1954-1958. Chesapeake Sci. 13: 87-100.
- Merriner, J.V. 1973. Assessment of the weakfish resource, a suggested management plan, and aspects of life history in North Carolina. Ph.D. Thesis, North Carolina State University, Raleigh, N.C., 201 p.

- Merriner, J.V. 1976. Aspects of the reproductive biology of the weakfish, Cynoscion regalis (Sciaenidae), in North Carolina. Fish. Bull., U.S. 74: 18-26.
- Nesbit, R.A. 1954. Weakfish migration in relation to its conservation. U.S. Fish. Wildlife Service Spec. Sci. Rep. Fish 115, 81 p.
- Nikolsky, G.V. 1963. The Ecology of Fishes. Academic Press, N.Y., 352 p.
- Patzner, R.A. and H. Adam. 1981. Changes in weight of the liver and the relationship to reproduction in the hagfish, Myxine glutinosa (Cyclostomata). J. Mar. Biol. Assoc. U.K. 61: 461-464.
- Paulson, A.C., and R.L. Smith. 1977. Latitudinal variation of Pacific herring fecundity. Trans. Am. Fish. Soc. 106(3): 244-247.
- Perlmutter, A., S.W. Miller, and J.C. Poole. 1956. The weakfish (Cynoscion regalis) in New York waters. N.Y. Fish Game J. 3: 1-43.
- Ricker, W.E. 1975. Computation and interpretation of biological statistics of fish populations. Bull. Fish. Res. Bd. Can. No. 191: 1-382.
- Schopka, S.A. 1971. Vergleichende Untersuchungen zur Fortpflanzungsrate bei Herings- und Kabeljaupopulationen (Clupea harengus L. und Gadus morhua L.). Ber. dt. wiss. komm. Meeresforsch. 22: 31-79. In: Hempel, G. 1979. Early life history of marine fish: the egg stage. Univ. Washington Press. 70 pp.
- Stearns, S.C. 1976. Life history tactics: A review of the ideas. Q. Rev. Biol. 51: 3-47.
- Thomas, D.L. 1971. The early life history and ecology of six species of drum (Sciaenidae) in the lower Delaware River, a brackish tidal estuary. Ichthyol. Assoc. Bull. 3, 247 p.
- Timashova, L.V. 1982. Seasonal changes in the structure of the liver in plaice Pleuronectes platessa. J. Ichthy. 21(2): 145-151.

- Welsh, W.W. and C.M. Breder. 1923. Contribution to the life histories of Sciaenidae of the eastern United States coast. Bull. U.S. Bur. Fish. 39: 141-201.
- Wenner, A.M. 1972. Sex ratio as a function of size in marine crustacea. Am. Nat. 106(949): 321-350.
- White, M.L. and M.E. Chittenden, Jr. 1977. Age determination reproduction and population dynamics of the Atlantic croaker, Micropogonias undulatus. Fish. Bull. 75(1): 109-123.
- Wilk, S.J. 1979. Biological and fisheries data of weakfish, Cynoscion regalis (Block and Schneider). NOAA Tech. Ser. Report No. 21.
- Wilk, S.J. 1981. A review of the fisheries for Atlantic croaker, spot, and weakfish, 1940-1979. In: H. Clepper (ed.), Marine Recreational Fisheries Symposium VI, Sportfishing Instit., Washington, D.C., 50 p.

TEXT FOOTNOTES

2,3,4

Shepherd G. and C.B. Grimes (In press). Geographic and historic variations in growth of weakfish (Cynoscion regalis) in the Middle Atlantic Bight. Fish. Bull., U.S.

5

National Ocean Survey, 1972. Surface water temperature and density; Atlantic Coast, North and South America. NOS publ. 31-1, 4th ed.

Table 1. Wilcoxon (Mann-Whitney) test comparing hepatosomatic indices between sexes of weakfish (Cynoscion regalis) from the New York Bight for June through November 1980 and May 1981. S = Wilcoxon test statistic, z = critical value.

1980	S	N	Z	Probability Level
June	622.0	50	5.081	0.0001 <sup>***</sup>
July	194.0	34	-1.767	0.0865
August	106.0	28	0.265	0.7928
September	274.0	37	-0.904	0.3658
October	252.0	35	0.017	0.9867
November	283.5	40	-1.215	0.2318
<hr/>				
<u>1981</u>				
May	332.0	48	-3.782	0.004 <sup>***</sup>

\*\*\* Highly Significant Differences

Tot  
Siz  
mat  
95%  
Int

Table 2. Length at maturity for weakfish  
(Cynoscion regalis) from 1980-83.

Length (cm)	FEMALE		MALE	
	N	% Mature	N	% Mature
18	15	0.0	16	0.0
19	7	0.0	11	0.0
20	8	12.5	11	9.1
21	11	18.2	5	40.0
22	8	37.5	9	33.3
23	13	53.8	16	31.3
24	11	45.5	12	50.0
25	13	46.2	12	75.0
26	14	71.4	15	86.7
27	17	88.2	14	71.4
28	10	80.0	20	65.0
29	17	82.4	22	63.6
30	19	100.0	17	82.4
31	18	88.9	14	85.7
32	16	93.8	20	85.0
33	10	90.0	17	94.1
34	14	85.7	15	100.0
35	7	85.7	3	100.0
36	10	80.0	8	100.0
37	9	77.8	12	100.0
38	15	100.0		
39	14	85.7		
40	13	92.3		
41	4	100.0		
42	10	100.0		
43	10	100.0		
Total	313		269	
Size at 50% mature	25.6		25.1	
95% Confidence Interval	24.4-26.8		24.0-26.2	

Table 3. Sex ratios (female:male) of weakfish  
(Cynoscion regalis) in the New York Bight.

Length <sup>a</sup>	Ratio	$\chi^2$
15	27:40	2.52
20	72:73	0.01
25	149:174	1.93
30	148:157	0.27
35	72:73	0.01
40	71:65	0.26
45	47:72	5.25*
50	35:40	0.33
55	34:60	7.19**
60	36:35	0.01
65	35:27	2.00
70	31:28	0.15
75	28:17	2.69
80	18:1	15.21***
85	4:0	4.0*
TOTAL	807:862	1.81

- \* significant difference  $P < 0.05$
- \*\* significant difference  $P < 0.01$
- \*\*\* significant difference  $P < 0.001$
- <sup>a</sup> midpoint of length interval (12.5 to 17.5, etc.)

Table 4. Difference between expected fecundity (based on  $\ln \text{ fec} = -16.322 + 4.659 \ln \text{ TL}$ ) and observed fecundity for 6 weakfish (Cynoscion regalis) collected 27 June 1981, in Gardiners Bay, N.Y.

FECUNDITY ESTIMATE			
Total Length	Expected	Observed	% Difference
590 mm	650,171	196,891	69.7
613 mm	777,718	93,579	88.0
615 mm	789,675	105,919	86.6
704 mm	1,487,262	438,909	70.5
737 mm	1,843,210	650,600	64.7
644 mm	979,893	313,054	<u>68.1</u>
		$\bar{x} =$	74.6 %

Table 5. Comparison of fecundity for weakfish (Cynoscion regalis) between Cape Hatteras (from Merriner 1976) and New York Bight.

FECUNDITY COMPARISONS				
Age	<u>Cape Hatteras*</u>		<u>New York Bight</u>	
	SL	Fecundity	TL	Fecundity
0	159	149,429		
1	225	391,688	203	4,593
2	286	762,258	323	39,978
3	357	1,410,550	479	250,685
4	421	2,229,220	578	601,544
5	562	4,969,940	638	953,027
6			677	1,256,487
7			701	1,477,902
8			728	1,762,445
9			758	2,127,278
10			763	2,193,448
CUMULATIVE FECUNDITY		9,913,085		10,667,390

\* Merriner, 1976

Table 6. Surface water temperatures (°C) for April-July  
in Plum Island, NY; Cape May, NJ; Gloucester Point,  
VA; and Beaufort, NC.<sup>5</sup>

	APRIL	MAY	JUNE	JULY
<u>PLUM ISLAND, NY</u>				
$\bar{X}$	5.6	9.7	14.6	18.8
Max	9.0	18.0	20.0	24.0
Min	2.0	4.0	9.0	15.0
$\bar{X}_{\max}$	7.7	13.4	17.7	21.4
$\bar{X}_{\min}$	3.3	6.4	11.4	16.3
<u>CAPE MAY, NJ</u>				
$\bar{X}$	10.2	14.5	20.0	22.6
Max	17.0	21.0	25.0	28.0
Min	6.0	9.0	12.0	18.0
$\bar{X}_{\max}$	14.6	18.9	23.8	25.6
$\bar{X}_{\min}$	7.0	10.8	15.7	19.9

Table 6 (continued). Surface water temperatures (°C) for April-July in Plum Island, NY; Cape May, NJ; Gloucester Point, VA; and Beaufort, NC.<sup>5</sup>

	APRIL	MAY	JUNE	JULY
<u>GLOUCESTER POINT, VA</u>				
$\bar{X}$	13.1	19.1	23.8	26.5
Max	21.0	26.0	30.0	32.0
Min	7.0	13.0	17.0	21.0
$\bar{X}_{\max}$	17.1	22.4	27.0	29.0
$\bar{X}_{\min}$	9.6	15.6	20.4	24.1
<u>BEAUFORT, NC</u>				
$\bar{X}$	17.2	21.4	25.2	27.4
Max	21.0	26.0	32.0	30.0
Min	14.0	14.0	18.0	24.0
$\bar{X}_{\max}$	20.3	24.2	28.3	29.2
$\bar{X}_{\min}$	14.3	17.5	21.9	25.8

## FIGURES

- Fig. 1      Sampling locations for weakfish (Cynoscion regalis) in the New York Bight and North Carolina site of Merriner (1976).
- Fig. 2      Changes in visual maturity stage of weakfish (Cynoscion regalis) in 1980-81 for the New York Bight. N indicated for each month.
- Fig. 3      Mean monthly gonosomatic indices  $\pm$  95% CI of weakfish (Cynoscion regalis) for 1980 in the New York Bight. N values at each point.
- Fig. 4      Monthly gonosomatic indices of individual weakfish (Cynoscion regalis) by sex for 1980. Spawning fish indicated by x.
- Fig. 5      Mean monthly hepatosomatic indices  $\pm$  95% CI of weakfish (Cynoscion regalis) for 1980 with N indicated at each point.
- Fig. 6      Monthly frequency of oocyte diameters for weakfish (Cynoscion regalis) in the New York Bight for 1980-81.
- Fig. 7      Fecundity as a function of length for weakfish (Cynoscion regalis) in New York Bight for 1981.

Fig. 1. Sampling locations for weakfish (Cynoscion regalis) in the New York Bight and North Carolina site of Merriner (1976).






Fig. 2. Changes in visual maturity stage of weakfish (Cynoscion regalis) in 1980-81 for the New York Bight. N indicated for each month.



 RECOVERING  
  IMMATURE  
  DEVELOPING  
  RIPE  
  SPENT

Fig. 3. Mean monthly gonosomatic indices  $\pm$  95% CI of weakfish (Cynoscion regalis) for 1980 in the New York Bight. N values at each point.




Fig. 4. Monthly gonosomatic indices of individual weakfish (Cynoscion regalis) by sex for 1980. Spawning fish indicated by x.




Fig. 5. Mean monthly hepatosomatic indices  $\pm$  95% CI  
of weakfish (Cynoscion regalis) for 1980 with  
N indicated at each point.




Fig. 6. Monthly frequency of oocyte diameters for weakfish  
(Cynoscion regalis) in the New York Bight for 1980-81.




Fig. 7. Fecundity as a function of length for weakfish  
(Cynoscion regalis) in New York Bight for 1981.

